



(12) **United States Patent**
Mundheim

(10) **Patent No.:** **US 9,409,801 B2**
(45) **Date of Patent:** **Aug. 9, 2016**

(54) **ALGINATE IMPROVEMENT**
(75) Inventor: **Atle Mundheim**, Omastrand (NO)
(73) Assignee: **SORBWATER TECHNOLOGY AS**,
Bergen (NO)
(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 490 days.

(21) Appl. No.: **13/063,416**
(22) PCT Filed: **Jan. 19, 2011**

(86) PCT No.: **PCT/NO2011/000023**
§ 371 (c)(1),
(2), (4) Date: **Mar. 10, 2011**

(87) PCT Pub. No.: **WO2011/090391**
PCT Pub. Date: **Jul. 28, 2011**

(65) **Prior Publication Data**
US 2011/0284471 A1 Nov. 24, 2011

(30) **Foreign Application Priority Data**
Jan. 19, 2010 (NO) 20100088

(51) **Int. Cl.**
B01D 21/01 (2006.01)
C02F 1/52 (2006.01)
C02F 1/68 (2006.01)
C02F 1/54 (2006.01)
C02F 1/66 (2006.01)
B03D 3/00 (2006.01)
C02F 1/00 (2006.01)

(52) **U.S. Cl.**
CPC **C02F 1/5272** (2013.01); **C02F 1/54**
(2013.01); **C02F 1/66** (2013.01); **C02F 2209/06**
(2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS
3,951,797 A 4/1976 Seely
5,520,819 A 5/1996 Asahi et al.
6,267,893 B1 * 7/2001 Luxemburg 210/723
6,372,145 B1 4/2002 Tarbet
7,727,404 B2 6/2010 Ylikangas et al.
2003/0089668 A1 * 5/2003 Moffett C02F 1/5236
210/725
2005/0098499 A1 * 5/2005 Hussain 210/639
2006/0163164 A1 * 7/2006 Pratt 210/724

(Continued)
FOREIGN PATENT DOCUMENTS
CN 1410372 A 4/2003
JP 10076274 3/1998
WO WO 2009052362 A2 * 4/2009 C08L 3/02

OTHER PUBLICATIONS
Nicholas M. Velings et al, Physico-Chemical Properties of Alginate
Gel Beads, (1995) 311-330.
(Continued)

Primary Examiner — Nam Nguyen
Assistant Examiner — Clare Perrin
(74) *Attorney, Agent, or Firm* — Flener IP Law; Zareefa B.
Flener

(57) **ABSTRACT**
A method is described use of hydrocolloids for removal of
organic and inorganic pollutants from a continuous stream of
liquid where chemical composition of the water inhibits the
reactants for linking/cross linking hydrocolloids. The method
is comprising the steps of enabling said reactants present in
the liquid stream and bring them to reaction with the hydro-
colloid already brought to attachment with pollutants, where
after flock formation allows for separation. Applications of
the method are also described.

12 Claims, 1 Drawing Sheet

Typical pH peak (pH vs buffering time in seconds)

