

[54] COMPOSITION FOR TREATMENT OF WOUNDS

[76] Inventor: Anthony N. Silvetti, 930 Ashland Ave., River Forest, Ill. 60305

[21] Appl. No.: 301,472

[22] Filed: Sep. 14, 1981

Related U.S. Application Data

[63] Continuation of Ser. No. 117,717, Feb. 19, 1980, abandoned.

[51] Int. Cl.³ A61K 31/70; A61K 33/26

[52] U.S. Cl. 424/147; 424/180; 424/319; 424/DIG. 13

[58] Field of Search 424/180, 319, 147

References Cited

U.S. PATENT DOCUMENTS

3,812,252 5/1974 Silvetti 424/180
4,105,783 8/1978 Yu et al. 426/283

FOREIGN PATENT DOCUMENTS

7513892 11/1975 Netherlands 424/280

OTHER PUBLICATIONS

Chemical Abstracts 92:4024h, 1980 (McKeehan et al.).
The Merck Index, 9th ed., 1976, p. 695.

American Drug Index, 1975, p. 228.

Biddle's Materia Medica & Therapeutics, p. 431, 1895.

Chemical Abstracts 80:13050t (1974) (Walser).

Br. J. of Surgery, vol. 63, pp. 427-430, 1976, Viljanto et al.

Methods of Tissue Culture, 3rd Ed. (1962), Hoeber Med. Div., Harper & Rowe, Parker, pp. 53-61.

Primary Examiner—Leonard Schenkman

Attorney, Agent, or Firm—Silverman, Cass & Singer

[57] ABSTRACT

Treatment of skin wounds with starch hydrolysate composition can be improved by irrigating the wound with a buffered salt solution having a pH in the range of approximately 6-7.8 prior to application of the starch hydrolysate composition. Additionally, the treatment can include further irrigation with amino acid solution. A preferred starch hydrolysate composition is compounded to include alphaketoglutaric acid or a salt thereof.

6 Claims, No Drawings