



[54] COMPOSITION AND METHODS FOR TRANSFECTING EUKARYOTIC CELLS

[75] Inventor: Joel A. Jessee, Mt. Airy, Md.
[73] Assignee: Life Technologies, Inc., Gaithersburg, Md.
[*] Notice: This patent is subject to a terminal disclaimer.
[21] Appl. No.: 08/746,896
[22] Filed: Nov. 18, 1996

Related U.S. Application Data

[63] Continuation of application No. 08/274,397, Jul. 12, 1994, Pat. No. 5,578,475, which is a continuation-in-part of application No. 08/090,290, Jul. 12, 1993, abandoned.
[51] Int. Cl.7 C12N 15/64; C12N 7/00; C12N 7/06
[52] U.S. Cl. 435/458; 435/235.1; 435/236; 435/456
[58] Field of Search 435/440, 235.1, 435/458, 236, 456; 424/184.1, 209.1, 211.1, 218.1

References Cited

U.S. PATENT DOCUMENTS

4,946,787 8/1990 Eppstein et al. 264/4.1
5,578,475 11/1996 Jessee 435/456

FOREIGN PATENT DOCUMENTS

WOA91/16024 10/1991 WIPO.
WO93/07282 4/1993 WIPO.
WO93/07283 4/1993 WIPO.
WO95/21259 10/1995 WIPO.

OTHER PUBLICATIONS

Kraaijeveld, C.A. et al. (1984), "The effect of Liposomal Charge on the Neutralizing Antibody Response Against Inactivated Encephalomyocarditis and Semliki Forest Viruses," Clin. Exp. Immunol. 56:509-514.
Neugebauer, J.M. (1990), "Detergents: An Overview," Methods Enzymol. 182:239-253.
Klappe, K. et al. (1986), "Parameters Affecting Fusion between Sendai Virus and Liposomes. Role of Viral Proteins, Liposome Composition, and pH," Biochemistry 25:8252-8260.
Eytan, G.D. (1982), "Use of Liposomes for Reconstitution of Biological Functions," Biochem. Biophys. Acta 694:185-202.
Sands, J.A. (1986), "Virucidal activity of cetyltrimethylammonium bromide below the critical micelle concentration," FEMS Microbiol. Lett. 36:261-263.
Walker, C. et al. (1992), "Cationic lipids direct a viral glycoprotein into the class I major histocompatibility complex antigen-presentation pathway," Proc. Natl. Acad. Sci. USA 89:7915-7918.

Konopka, K. et al., (1991), "Enhancement of human immunodeficiency virus type 1 infection by cationic liposomes: the role of CD4, serum and liposome-cell interactions," J. Gen. Virol. 72:2685-2696.
Lapidot, M. and Loyter, A. (1990), "Fusion-Mediated Microinjection of Liposome-Enclosed DNA into Cultured Cells with the Aid of Influenza Virus Glycoproteins," Exp. Cell Res. 189:241-246.
Gould-Fogerite, S. et al. (1989), "Chimerasome-mediated gene transfer in vitro and in vivo," Gene 84:429-438.
Kaneda et al., (1987), "The Improved Efficient Method for Introducing Macromolecules into Cells Using HFM (Sendai virus) Liposomes with Gangliosides," Exp. Cell Res. 173:56-69.
Kaneda, Y. et al. (1989), "Introduction and Expression of the Human Insulin Gene in Adult Rat Liver," J. Biol. Chem. 264:12126-12129.
Väänänen, P. and Kääriäinen, L. (1980), "Fusion and Haemolysis of Erythrocytes caused by Three Togaviruses: Semliki Foest, Sinbis and Rubella," J. Gen. Virol. 46:467-475.
Young, J.D.-E. et al. (1983), "Interaction of Enveloped Viruses with Planar Bilayer Membranes: Observations on Sendai, Influenza, Vesicular Stomatitis, and Semliki Forest Viruses," Virology 128:186-194.
Marsh, M. et al. (1983), "Interactions of Semliki Forest Virus Spike Glycoprotein Rosettes and Vesicles with Cultured Cells," J. Cell. Biol. 96:455-461.
Scheule, R.K. (1986), "Novel Preparation of Functional Sindbis Virosomes," Biochemistry 25:4223-4232.
Ciccarone, V. et al. (1993), "Cationic liposome-mediated transfection of Eukaryotic cells high efficiency nucleic acid delivery with lipofectin lipofectace, and lipofectamine reagents," in FASEB Journal, Abstracts, American Society for Biochemistry and Molecular Biology and Division of Biological Chemistry—American Chemical Society, joint meeting, San Diego, CA, May 30-Jun. 3, 1993, Abstract No. 454, p. A1131.
"Transfection Reagent," Genet. Eng. News (Jun. 15, 1993), p. 12, col. 4.
Phalen, T. and Kielian, M. (1991), "Cholesterol Is Required for Infection by Semliki Forest Virus," J. Cell Biol. 112:615-623.

(List continued on next page.)

Primary Examiner—Terry McKelvey
Attorney, Agent, or Firm—Greenlee, Winner & Sullivan, P.C.

ABSTRACT

[57] The present invention provides compositions and methods for transfecting eukaryotic cells which comprise a nucleic acid, a cationic lipid capable of forming a complex with said nucleic acid, and a viral agent which is an active or inactive enveloped virus or a component of an enveloped virus. Particular viral agents include alpharviruses and components thereof.

97 Claims, No Drawings