



US005597607A

United States Patent [19]

[11] Patent Number: **5,597,607**

Samoto et al.

[45] Date of Patent: **Jan. 28, 1997**

[54] **PROCESS FOR PREPARING FRACTIONATED SOYBEAN PROTEINS AND FOODS USING THE SAME**

[75] Inventors: **Masahiko Samoto; Takeshi Akasaka**, both of Tsukuba-gun; **Hiroyuki Mori**, Abiko, all of Japan

[73] Assignee: **Fuji Oil Company, Limited**, Osaka-fu, Japan

[21] Appl. No.: **364,102**

[22] Filed: **Dec. 27, 1994**

[30] **Foreign Application Priority Data**

Jan. 7, 1994	[JP]	Japan	6-000620
Nov. 22, 1994	[JP]	Japan	6-288425

[51] Int. Cl.⁶ **A23J 1/14**; A23J 3/16; A23L 1/20

[52] U.S. Cl. **426/656**; 530/378

[58] Field of Search 426/656; 530/378

[56] **References Cited**

FOREIGN PATENT DOCUMENTS

0072094 2/1983 European Pat. Off. .

OTHER PUBLICATIONS

Ogawa et al J. Nutri. Sci Vitaminol., 37, 555-565, (1991).
Thanh et al J. Agric. Food Chem. vol. 24, No. 6 pp. 1117-1121 (1976).

Sato et al Phytochemistry, vol. 23, No. 8 pp. 1523-1526, 1984.

Abstract of JP-52-035739, NODA Inst Sci Res, "Soyabean 7S Protein Production by Dispersing Soyaben GLobulin in a Solution Containing Sodium Chloride or Potassium Chloride", Sep. 10, 1977.

Abstract of JP-5-043597, Nagano et al, "Method for Fractioning 7S Protein", Feb. 23, 1993.

Abstract of JP-55-124457, Koshiyama et al., "Preparation of 7S Protein", Sep. 25, 1980.

Anderson et al., "Extraction of Soybean Meal Proteins with Salt Solutions at pH 4.5", J. Agr. Food Chem, vol. 21, No. 2, pp. 251-254, 1973.

Iwabuchi et al., "Determination of GLy cinin and β -Conglycinin in Soybean Proteins by Immunological Methods", J. Agr. Food Chem. vol. 35, No. 2, pp. 200-205, 1987.

Primary Examiner—Esther M. Kepplinger

Assistant Examiner—Choon P. Koh

Attorney, Agent, or Firm—Wenderoth, Lind & Ponack

[57] **ABSTRACT**

A novel, efficient and simple process for preparing low-allergenic fractionated soybean proteins by treating soybean proteins under acidic conditions with an aqueous solution in which a salt is dissolved to selectively concentrate Gly m Bd 30k in a precipitation fraction and collecting a supernatant fraction.

11 Claims, 3 Drawing Sheets