

**PROTEINS WITH REPETITIVE
BACTERIAL-IG-LIKE (BIG) DOMAINS
PRESENT IN *LEPTOSPIRA* SPECIES**

CROSS REFERENCE TO RELATED
APPLICATIONS

This is a divisional of U.S. application Ser. No. 14/819,045, now U.S. Pat. No. 9,346,858, filed Aug. 5, 2015; which is a divisional of U.S. application Ser. No. 14/281,580, filed May 19, 2014, now U.S. Pat. No. 9,133,250, issued Sep. 15, 2015; which is a divisional of U.S. application Ser. No. 13/869,660, filed Apr. 24, 2013, now U.S. Pat. No. 8,802,835, issued Aug. 12, 2014; which is a divisional of U.S. application Ser. No. 13/525,157, filed Jun. 15, 2012, now U.S. Pat. No. 8,445,658, issued May 21, 2013; which is a divisional of U.S. application Ser. No. 13/359,354, filed Jan. 26, 2012, now U.S. Pat. No. 8,216,594, issued Jul. 10, 2012; which is a divisional of U.S. application Ser. No. 13/216,214, filed Aug. 23, 2011, now U.S. Pat. No. 8,124,110, issued Feb. 28, 2012; which is a divisional of U.S. application Ser. No. 13/078,879, filed Apr. 1, 2011, now U.S. Pat. No. 8,021,673, issued Sep. 20, 2011; which is a divisional of U.S. application Ser. No. 12/728,177, filed Mar. 19, 2010, now U.S. Pat. No. 7,935,357, issued May 3, 2011; which is a divisional of application Ser. No. 11/332,464, filed Jan. 17, 2006, now U.S. Pat. No. 7,718,183, issued May 18, 2010; which is a divisional of U.S. application Ser. No. 11/005,565, filed Dec. 7, 2004 (abandoned); which is a divisional of U.S. application Ser. No. 10/147,299, filed May 17, 2002 (abandoned). The entire contents of each of the earlier applications is hereby incorporated by reference.

ACKNOWLEDGEMENT OF GOVERNMENT
SUPPORT

This invention was made with Government support under Grant Nos. AI001605, AI034431, HL051967, and TW000905 awarded by the National Institutes of Health. The Government has certain rights in this invention.

INCORPORATION-BY-REFERENCE OF
MATERIAL SUBMITTED AS AN ASCII TEXT
FILE

A Sequence Listing is submitted herewith as an ASCII compliant text file named "Sequence_Listing.txt", created on Apr. 20, 2016, and having a size of 78,445 bytes, as permitted under 37 CFR 1.821(c). The material in the aforementioned file is hereby incorporated by reference in its entirety.

FIELD

The invention relates to three isolated DNA molecules that encode for proteins, BigL1, BigL2 and BigL3, in the *Leptospira* sp bacterium which have repetitive Bacterial-Ig-like (Big) domains and their use in diagnostic, therapeutic and vaccine applications. According to the present invention, the isolated molecules encoding for BigL1, BigL2 and BigL3 proteins are used for the diagnosis and prevention of infection with *Leptospira* species that are capable of producing disease in humans and other mammals, including those of veterinary importance.

BACKGROUND

Spirochetes are motile, helically shaped bacteria and include three genera, *Leptospira*, *Borrelia* and *Treponema*,

which are pathogens of humans and other animals. *Borrelia* and *Treponema* are the causative agents of diseases that include Lyme disease, relapsing fever, syphilis and yaws. *Leptospira* consists of a genetically diverse group of eight pathogenic and four non-pathogenic, saprophytic species (1, 2). Leptospire are also classified according to serovar status—more than 200 pathogenic serovars have been identified. Structural heterogeneity in lipopolysaccharide moieties appears to be the basis for the large degree of antigenic variation observed among serovars (1, 2).

Leptospirosis is a zoonotic disease; transmission to humans occurs through contact with domestic or wild animal reservoirs or an environment contaminated by their urine. Infection produces a wide spectrum of clinical manifestations. The early-phase of illness is characterized by fever, chills, headache and severe myalgias. Disease progresses in 5 to 15% of the clinical infections to produce severe multisystem complications such as jaundice, renal insufficiency and hemorrhagic manifestations (1-4). Severe leptospirosis is associated with mortality rates of 5-40%.

Leptospirosis has a world-wide distribution. Because of the large spectrum of animal species that serve as reservoirs, it is considered to be the most widespread zoonotic disease (1). Leptospirosis is traditionally an important occupational disease among risk groups such as military personnel, farmers, miners, sewage and refuse removal workers, veterinarians and abattoir workers (1-3). However, new patterns of disease transmission have emerged recently that emphasize the growing importance of leptospirosis as a public health problem. In developed countries, leptospirosis has become the cause of outbreaks associated with recreational activities (1) and sporting events (1, 4, 5). In Brazil and other developing countries, underlying conditions of poverty have produced large urban epidemics of leptospirosis associated with high mortality (4, 5).

In addition to its public health impact, leptospirosis is a major economic burden as the cause of disease in livestock and domestic animals (2). Leptospirosis produces abortions, stillbirths, infertility, failure to thrive, reduced milk production and death in animals such as cows, pigs, sheep, goats, horses and dogs and induces chronic infection and shedding of pathogenic leptospire in livestock (2) and therefore represents an additional source of economic loss for the animal husbandry industry because of current international and national quarantine regulations.

The control of human and animal leptospirosis is hindered by the current lack of adequate diagnostic tools. The standard serologic test, the microscopic agglutination test (MAT), is inadequate for rapid case identification since it can only be performed in few reference laboratories and requires analyses of paired sera to achieve sufficient sensitivity (1, 2). Dependence upon the MAT results in delays in establishing the cause of outbreaks as seen in several investigations (1, 2). Enzyme-linked immunosorbent assays (ELISA), and other rapid serologic tests based on whole-cell leptospiral antigen preparations have been developed for use as an alternative method to screen for leptospiral infection, although the MAT is still required for case confirmation (1, 2). Recombinant antigen-based serologic tests are widely used in screening for spirochetal infections such as Lyme disease and syphilis, but the use of recombinant proteins for serodiagnosis of leptospirosis has not been widely investigated. Recently, a recombinant flagellar-antigen immunocapture assay was described for serodiagnosis of bovine leptospirosis (6). A recombinant heat shock protein, Hsp58, showed a high degree of ELISA reactivity with serum samples from a small number of human cases (7). However,