United States Patent [19]

Sjöquist

[11] **3,995,018**

[45] **Nov. 30, 1976**

[54] **METHOD OF BINDING IMMUNOGLOBULIN EMPLOYING A POLYPEPTIDE FROM MICROORGANISMS**

[75] Inventor: **John Axel Sjöquist,** Uppsala, Sweden

[73] Assignee: **Pharmacia Aktiebolag,** Uppsala, Sweden

[22] Filed: **Oct. 25, 1973**

[21] Appl. No.: **409,572**

[30] **Foreign Application Priority Data**

Nov. 6, 1972 Sweden 14330/72
Feb. 8, 1973 Sweden 7301779

[52] **U.S. Cl.** ..... **424/1.5;** 260/112 R; 424/7; 424/12; 424/88

[51] **Int. Cl.**[2] **G01N 33/00;** C07G 7/00; A61K 39/00; A61K 43/00

[58] **Field of Search** 424/1, 12, 88, 359

[56] **References Cited**

UNITED STATES PATENTS

3,555,143 1/1971 Axen et al. 424/1
3,788,948 1/1974 Kagedal et al. ..... 424/1 X
3,790,663 2/1974 Garrison et al. 424/12
3,850,798 11/1974 Sjoquist ..... 210/31 C

FOREIGN PATENTS OR APPLICATIONS

7,018,879 6/1970 Japan 424/12

OTHER PUBLICATIONS

Bambardieri et al., Proc. Soc. Exp. Biol. Med., vol. 133, No. 4, Apr. 1970, pp. 1366–1369.

Primary Examiner—Benjamin R. Padgett
Assistant Examiner—Christine M. Nucker
Attorney, Agent, or Firm—Fred Philpitt

[57] **ABSTRACT**

A method of binding, in the presence of a liquid, at least one immunoglobulin or its free Fc-fragment, the immunoglobulin being in free form or with its Fab-part bound to an antigen to which is optionally in turn bound one or more groups or substances, to a polymer insoluble in said liquid with the assistance of a substance attached to the polymer. The substance used is at least one polypeptide from microorganisms to which polypeptide the Fc-part of the immunoglobulin or its free Fc-fragment can bind itself.

11 Claims, No Drawings