# United States Patent [19]

**Eisinger**

[11] **4,299,819**

[45] **Nov. 10, 1981**


[54] **PROCESS FOR TREATING BURN VICTIMS**


[75] Inventor: **Magdalena G. Eisinger,** Demarest, N.J.

[73] Assignee: **Sloan-Kettering Institute for Cancer Research,** New York, N.Y.



[21] Appl. No.: **75,375**

[22] Filed: **Sep. 13, 1979**

**Related U.S. Application Data**

[62] Division of Ser. No. 749, Jan. 2, 1979.

[51] **Int. Cl.**[3] ..... **A61K 35/12;** A01N 1/02

[52] **U.S. Cl.** ..... **424/95;** 435/1; 435/241

[58] **Field of Search** ..... 424/95, 366; 435/1, 435/240, 241



[56] **References Cited**

U.S. PATENT DOCUMENTS

4,016,036 4/1977 Green et al. ..... 435/241

OTHER PUBLICATIONS

Kitano et al.-Biochem. of Cutaneous Epidermal Differentiation, Ed. by Seiji, (1977), pp. 319-335.

Rheinwald et al.-Cell, vol. 6, Nov. 1975, pp. 313-334.

Eagle-Science, vol. 174, Oct. 29, 1971, pp. 501-503.

Eagle-J. Cellular Physiology, vol. 82, No. 1, Aug. 1973, pp. 1-7.

Willmer-Cells & Tissue in Culture, vol. 1 (1965), pp. 24 & 25.



*Primary Examiner*—Sam Rosen

*Attorney, Agent, or Firm*—Sprung, Felfe, Horn, Lynch & Kramer



[57] **ABSTRACT**

Burn victims are treated with human epidermis cells grown in tissue culture by separating the epidermis in human skin from the dermis, dissociating the epidermis into epidermal cells, and growing the epidermal cells in a tissue culture medium having a pH of from about 5.6 to 5.8.


**13 Claims, No Drawings**