United States Patent [19]

Bayer et al.

[11] **3,960,830**

[45] **June 1, 1976**

[54] **POLYALKYLENE GLYCOLS USED FOR THE PREPARATION OF PEPTIDES**

[75] Inventors: **Ernst Bayer; Manfred Mutter,** both of Tubingen, Germany

[73] Assignee: **Hoechst Aktiengesellschaft,** Frankfurt am Main, Germany

[22] Filed: **Dec. 4, 1974**

[21] Appl. No.: **529,519**

[30] **Foreign Application Priority Data**

Dec. 6, 1973 Germany 2360794

[52] **U.S. Cl.** ..... **260/112.5 R**

[51] **Int. Cl.**[2] **C07C 103/52**

[58] **Field of Search** 260/112.5

[56] **References Cited**

UNITED STATES PATENTS

3,502,545 3/1970 Westman et al. 260/112.5

3,645,852 2/1972 Axen et al. 260/112.5

OTHER PUBLICATIONS

Bayer et al.: Nature, 237, pp. 512–513, (1972).

Primary Examiner—Lewis Gotts
Assistant Examiner—Reginald J. Suyat
Attorney, Agent, or Firm—Curtis, Morris & Safford

[57] **ABSTRACT**

A process for the synthesis of peptides in a liquid phase wherein the peptide product, and intermediates in the peptide synthesis, are bound to a polyalkylene glycol polymer and wherein the polymer-bound peptide is separated from the liquid phase, e.g. for purposes of isolation or purification, by crystallizing the polymer-bound peptide.

2 Claims, No Drawings