U.S. PATENT DOCUMENTS

5,674,908	A	10/1997	Haces et al.	514/642
5,714,166	A	2/1998	Tomalia et al.	
5,736,387	A	4/1998	Paul et al.	435/320.1
5,736,392	A	4/1998	Hawley-Nelson et al.	
5,759,805	A	6/1998	Feldhaus et al.	435/69.1
5,773,527	A	6/1998	Tomalia et al.	525/417
5,795,587	A	8/1998	Gao et al.	424/450
5,837,533	A	11/1998	Boutin	435/320.1
5,854,224	A	12/1998	Lockett et al.	514/44
5,906,922	A	5/1999	Whittaker et al.	435/69.1

OTHER PUBLICATIONS

Curiel, D.T. et al., "Gene transfer to respiratory epithlial cells via the receptor–mediated endocytosis pathway," (1992) *Am. J. Respir. Cell. Mol. Biol.* 6:247–252.

Demeneix, B.A. et al., "Gene transfer into intact vertebrate embryos," (1991) *Int. J. Dev. Biol.* 35:481–484.

Dwarki, V.J., "Cationic liposome–mediated RNA transfection," (1993) *Methods in Enzymology* 217:644–654.

Finlay, D.R. et al., "Nuclear transport in vitro," (1989) *J. Cell Sci. Suppl. 11* p. 225–242.

Gao, X and Huang, L., "A novel cationic liposome reagent for efficient transfection of mammalian cells," (1991) *Biochem. and Biophys. Res. Communications* 179(1):280–285.

Garcia–Bustos, J. et al., "Nuclear protein localization," (1991) *Biochimica et Biophysica Acta* 1071:83–101.

Goldfarb, D.S. et al., "Synthetic peptides as nuclear localization signals," (1986) *Nature* 322:641–644.

Goldfarb, D.S. and Michaud, N., "Pathways for the nuclear transport of proteins and RNAs," (1991) *Trends in Cell Biology* 1:20–24.

Huckett, B. et al., "Evidence for targeted gene transfer by receptor–mediated endocytosis," (1990) *Biochem. Pharmacology* 40(2):253–263.

Kaneda, Y. et al., "Increased expression of DNA cointroduced with nuclear protein in adult rat liver," (1989) *Science* 243:375–378.

Karlsson, S. et al., "Transfer of genes into hematopoietic cells using recombinant DNA viruses," (1985) *Proc. Natl. Acad. Sci. USA* 82:158–162.

Lanford, R.E. et al., "Comparison of diverse transport signals in synthetic peptide–induced nuclear transport," (1990) *Exp. Cell Res.* 186:32–38.

Ledley, F.D., "Clinical considerations in the design of protocols for somatic gene therapy," (1991) *Human Gene Therapy* 2:77–83.

Legendre, J.–Y. and Szoka, F.C., Jr., "Delivery of plasmid DNA into mammalian cell lines using pH–sensitivie liposomes: Comparison with cationic liposomes," (1992) *Pharm. Res.* 9(10):1235–1242.

Legendre, J.–Y. and Szoka, F.C., Jr., "Cyclic amphipathic peptide–DNA complexes mediate high–efficiency transfection of adherent mammalian cells," (1993) *Proc. Natl. Acad. Sci. USA* 90:893–897.

Loyter, A. et al., "Mechanisms of DNA uptake by mammalian cells: Fate of exogenously added DNA monitored by the use of fluorescent dyes," (1982) *Proc. Natl. Acad. Sci. USA* 79:422–426.

Malone, R.W. et al., "Cationic liposome–mediated RNA transfection," (1989) *Proc. Natl. Acad. Sci. USA* 86:6077–6081.

Parente, R.A. et al., "Association of a pH–sensitive peptide with membrane vesicles: Role of amino acid sequence," (1990) *Biochemistry* 29:8713–8719.

Parente, R.A. et al., "Mechanism of leakage of phospholipid vesicle contents induced by the peptide GALA," (1990) *Biochemistry* 29:8720–8728.

Poste, G. et al., "Lipid vesicles as carriers for introducing biologically active materials into cells," (1976) *Methods in Cell Biology* 14:33–71.

Rosenkranz, A.A. et al., "Receptor–mediated endocytosis and nuclear transport of a tranfecting DNA construct," (1992) *Exp. Cell Res.* 199:323–329.

Schmid, N. and Behr, J.–P., "Location of spermine and other polyamines on DNA as revealed by photoaffinity cleavage wiht polyaminobenzenediazonium salts," (1991) *Biochemistry* 30:4357–4361.

Silver, P.A., "How proteins enter the nucleus," (1991) *Cell* 64:489–497.

Smull, C.E. and Ludwig, E.H., "Enhancement of the plaque– forming capacity of poliovirus ribonucleic acid with basic proteins," (1962) *J. Bacteriology* 84:1035–1040.

Sugawa, H. et al., "Large macromolecules can be introduced into cultured mammalian cells using erythrocyte membrane vesicles," (1985) *Exp. Cell Res.* 159:410–418.

Trubetskoy, V.S. et al., "Use of N–terminal modified poly (L–lysine)–antibody conjugate as a carrier for targeted gene delivery in mouse lung endothelial cells," (1992) *Bioconjugate Chem.* 3:323–327.

van Zee, K. et al., "A hydrophobic protein sequence can override a nuclear localization signal independently of protein context," (1991) *Mol. and Cellular Biol.* 11(10):5137–5146.

Wagner, E. et al., "DNA–binding transferrin conjugates as functional gene–delivery agents: Synthesis by linkage of polylysine or ethidium homodimer to the transferrin carbohydrate moiety," (1991) *Bioconjugate Chem.* 2:226–231.

Wagner, E. et al., "Transferrin–polycation conjugates as carriers for DNA uptake into cells," (1990 *Proc. Natl. Acad. Sci. USA* 87:3410–3414.

Wilson, J.M. et al., "Hepatocyte–directed gene transfer in vivo leads to transient improvement of hypercholesterolemia in low density lipoprotein receptor–deficient rabbits," (1992) *J. Biol. Chem.* 267(2):963–967.

Wolff, J.A. et al., "Direct gene transfer into mouse muscle in vivo," (1990) *Science* 247:1465–1468.

Wu, C.H. et al., "Targeting genes: Delivery and persistent expression of a foreign gene driven by mammalian regulatory elements in vivo," (1989) *J. Biol. Chem.* 265(29):16985–16987.

Wu, G.Y. et al., "Receptor–mediated gene delivery in vivo: Partial correction of genetic analbuminemia in nagase rats," (1991) *J. Biol. Chem.* 266(22):14338–14342.

Wu, G.Y. and Wu, C.H., "Receptor–mediated gene delivery and expression in vivo," (1988) *J. Biol. Chem.* 263(29):14621–14624.

Wu, G.Y. and Wu, C.H., "Evidence for targeted gene delivery to Hep G2 hepatoma cells in vitro," (1988) *Biochemistry* 27:887–892.

Yagi, K. et al., "Incorporation of histone into liposomes increases the efficiency of liposome–mediated gene transfer," (1991) *J. Clin. Biochem. Nutr.* 10:21–25.