OTHER PUBLICATIONS

Allred, J.C., et al., High–sensitivity atomic magnetometer unaffected by spin–exchange relaxation, *Phys. Rev. Lett.* (2002) 89:130801–1–130801–4.

Bison, G., et al., A laser–pumped magnetometer for the mapping of human cardiomagnetic fields, *Appl. Phys. B.* (2003) 76:325–328.

Bison, G., et al., Dynamical mapping of the human cardiomagnetic field with a room–temperature, laser–optical sensor, Opt. Expr. (2003) 11:904–909.

Budker, D., et al., Resonant nonlinear magneto–optical effects in atoms, *Rev. Mod. Phys.* (2002) 74:1153–1201.

Budker, D. et al., Nonliner Magneto–optic Effects with Ultranarrow Widths, *Phys. Rev. Lett.* (1998) 81:5788–5791.

Budker, D., et al., Sensitive magnetometry based on non–linear magneto–optical rotation, *Phys. Rev. A* (2000) 62:043403–1–043403–7.

Clem, T.R., Superconducting Magnetic Gradiometers For Underwater Target Detection, *Nav. Eng. J.* (1998) 110:139–149.

Del Gratta C, et al., Magnetoencephalography—a noninvasive brain imaging method with 1 ms time resolution, *Rep. Prog. Phys.* (2001) 64:1759–1814.

Drung, D., et al., Improved direct–coupled cd SQUID read–out electronics with automatic bias voltage tuning, *IEEE T. Appl. Supercon.* (2001) 11:880–883.

Greenberg, Ya.S., Application of superconducting quantum interference devices to nuclear magnetic resonance, *Rev. Mod. Phys.* (1998) 70:175–222.

Hämäläinen M. et al., Magnetoencephalography–theory, instrumentation, and applications to non–invasive studies on the working human brain, *Rev. Mod. Phys.* (1993) 65:413–497 (Abstract and contents only).

Happer, W., Optical Pumping, *Rev. Mod. Phys.* 1972, 44:169–249 (Abstract and contents only).

Happer, W., et al., Effect of rapid spin exchange on the magnetic–resonance spectrum of alkali vapors, *Phys. Rev. A* (1977) 16:1877–1991.

Happer W. et al., Spin–Exchange Shift and Narrowing of Magnetic Resonane Lines inOptically Pumped Akali Vapors, Phys. Rev. Lett. (1973) 31:273–276.

Harry, G.M., et al., Two–stage superconducting–quantum–interference–device amplifier in a high–Q gravitational wave transducer, *Appl. Phys. Lett.* (2000) 76:1446–1448.

Kelha, V.O., et al., Design, Construction and Performance of a large–volume magnetic shield, IEEE Trans. Magn. (1982) 18:260–270.

Kirschvink, J.L., et al., Paleomagnetic evidence of a low–temperature origin of carbonate in the Martian meteorite ALH84001, *Science* (1997) 275:1629–1633.

Kominis, I. K. et al., A subfemtotesia multichannel atomic magnetometer, Nature (2003) 422:596–599.

McDermott, R., et al., Liquid–state NMR and scalar couplings in microtesia magnetic fields, *Science* (2002) 295:2247–2249.

Murthy, S. A., et al, New Limits on the Electron Electric Dipole Moment from Cesium, Phys. Rev. Lett. (1989) 63:965–968.

Nenone, J., et al., Thermal noise in biomagnetic measurements, *Rev. Sci. Instr.* (1996) 67:2397–2405.

Nenonen, J., et al., Thermal Noise in a Magnetically Shielded Room, in Biomagnetism '87 , Ed. K. Atsumi et al., Denki University Press, Tokyo, 1988), p. 426–429.

Oukhanski, N., et al., Low–drift broadband directly coupled dc SQUID read–out electronics, *Physica C* (2002) 368:166–170.

Rodriguez, E., et al., Perception's shadow: long–distance synchronization of human brain activity, *Nature* (1999) 397:430–433.

Tralshawala, N., et al., Practical SQUID instrument for non–destructive testing, *Appl. Phys. Lett.* (1997) 71:1573–1575.

Ts'o, D.Y., et al., Functional organisation of primate visual cortex revealed by high–resolution optical imaging, *Science* (1990) 249:417–420.

Tsuei, C.C. et al., Phase–sensitive evidence for d–wave pairing symmetry in electron–doped cuprate superconductors, *Phys. Rev. Lett.* (2000) 85:182–185.

Varpula, T., et al., *J. App. Phys.* (1984) 55:4015–4021.

Zimmerman, J.E., et al., Design and operation of stable RF–biased superconducting point–contact quantum devices, and a note on properties of perfectly clean metal contacts. *J. Appl. Phys.* (1970) 41, 1572–1580.

* cited by examiner