

[54] KINETIC METHOD FOR DETERMINATION OF GLUCOSE CONCENTRATIONS WITH GLUCOSE DEHYDROGENASE

[75] Inventors: Kenneth J. Pierre; Ker-Kong Tung, both of Vista, Calif.

[73] Assignee: Beckman Instruments, Inc., Fullerton, Calif.

[21] Appl. No.: 791,825

[22] Filed: Apr. 28, 1977

[51] Int. Cl.² G01N 31/14

[52] U.S. Cl. 195/103.5 C

[58] Field of Search 195/103.5 C, 103.5 R

[56] References Cited

U.S. PATENT DOCUMENTS

3,964,974 6/1972 Banauch et al. 195/103.5 C

OTHER PUBLICATIONS

Banauch et al. *Z. Klin. Chem. Klin. Biochem*, vol. 3, pp. 101-107 (1975).

Lutz et al., *Clin. Chem.* 21: 1372-1377 (1975).

Lutz, *Clin. Chem.* 22: 929 (1976).

Rindfrey et al., *Z. Anal. Chem.* 279: 196 (1976).

Keller et al., *J. Clin. Chem. Clin. Biochem* 14: 27-30 (1976).

Pauley et al., *Hoppe-Seyler's Z. Physiol. Chem.* 356:1613-1623 (1975).

Primary Examiner—Lionel M. Shapiro

Attorney, Agent, or Firm—R. J. Steinmeyer; R. S.

Frieman

[57]

ABSTRACT

The range of linearity of the kinetic method for the determination of glucose concentrations using a reagent of the type comprising a buffer, a pyridine coenzyme, and glucose dehydrogenase, is significantly increased when the buffer employed in the reagent has a pH of from about 7.9 to about 9.0.

14 Claims, 1 Drawing Figure