



US005128360A

United States Patent [19]

[11] Patent Number: **5,128,360**

Cerami et al.

[45] Date of Patent: **Jul. 7, 1992**

[54] **METHOD AND AGENTS FOR INHIBITING PROTEIN AGING**

[75] Inventors: **Anthony Cerami**, Shelter Island, N.Y.; **Peter C. Ulrich**, Tenafly, N.J.; **Michael Brownlee**, New York, N.Y.

[73] Assignee: **The Rockefeller University**, New York, N.Y.

[21] Appl. No.: **481,869**

[22] Filed: **Feb. 20, 1990**

Related U.S. Application Data

[60] Continuation-in-part of Ser. No. 220,504, Jul. 18, 1988, abandoned, which is a division of Ser. No. 798,032, Nov. 14, 1985, Pat. No. 4,758,583, which is a division of Ser. No. 590,820, Mar. 19, 1984, Pat. No. 4,665,192.

[51] Int. Cl.⁵ **A61K 31/415; A61K 31/155**

[52] U.S. Cl. **514/400; 514/634; 514/866; 514/632**

[58] Field of Search **514/634, 866, 400, 632**

[56] **References Cited**

U.S. PATENT DOCUMENTS

| | | | |
|-----------|--------|----------------------|---------|
| 3,637,850 | 1/1972 | Houlihan et al. | 260/564 |
| 3,746,764 | 7/1973 | Nordmann et al. | 260/564 |
| 4,758,583 | 7/1988 | Cerami et al. | 514/399 |
| 4,908,446 | 3/1990 | Ulrich et al. | 540/553 |

FOREIGN PATENT DOCUMENTS

| | | |
|---------|---------|----------------------|
| 134099 | 11/1969 | Czechoslovakia . |
| 0111211 | 6/1984 | European Pat. Off. . |

OTHER PUBLICATIONS

Merck Index (Tenth Edition) 1983, p. 66 #444.

Brownlee et al., *Annals of Internal Medicine*, 101, pp. 527-537 (1984).

Dabrowski et al., *Acta Physiol. Pol.*, 34(1), pp. 91-97, (1983).

Diabetes, 35 (supp. 1) p. 42A (1986).

CA 113:34670k, Lewis et al.

CA 111:70141f, Brownlee.

CA 111:37368n, Nicholls et al.

CA 110:643b, Khatami et al.

CA 112:42594h, Onada et al.

CA 111:70766p, Oimomi et al.

CA 105:72441d, Brownlee et al.

Primary Examiner—Frederick E. Waddell

Assistant Examiner—Kimberly R. Jordan

Attorney, Agent, or Firm—Klauber & Jackson

[57] **ABSTRACT**

The present invention relates to compositions and methods for inhibiting protein aging. Accordingly, a composition is disclosed which comprises an agent or compound capable of inhibiting the formation of advanced glycosylation end products of target proteins by reacting with the carbonyl moiety of the early glycosylation product of such target proteins formed by their initial glycosylation. Suitable agents may contain an active nitrogen-containing group, such as a hydrazine group. Particular agents comprise aminoguanidine, α -hydrazinohistidine and mixtures thereof. The method comprises contacting the target protein with the composition. Both industrial and therapeutic applications for the invention are envisioned, as food spoilage and animal protein aging can be treated.

28 Claims, 5 Drawing Sheets