TREATMENT WITH ANTI-ERBB2 ANTIBODIES

This application is a continuation of U.S. application Ser. No. 11/780,640, filed Jul. 20, 2007, now U.S. Pat. No. 8,075,892 which application is a divisional of U.S. application Ser. No. 10/909,998, filed Aug. 2, 2004, which is a continuation of U.S. application Ser. No. 09/209,023, filed Dec. 10, 1998 (now abandoned) which claims priority under 35 U.S.C. Section 119(e) and the benefit of U.S. Provisional Application Ser. No. 60/069,346, filed Dec. 12, 1997, the entire disclosures of which are hereby incorporated by reference.

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Sep. 30, 2011, is named GNE0329C.txt and is 5,761 bytes in size.

FIELD OF THE INVENTION

The present invention concerns the treatment of disorders characterized by the overexpression of ErbB2. More specifically, the invention concerns the treatment of human patients susceptible to or diagnosed with cancer overexpressing ErbB2 with a combination of an anti-ErbB2 antibody and a chemotherapeutic agent other than an anthracycline, e.g. doxorubicin or epirubicin. The invention further provides a method of treating cancer in a human patient comprising administering effective amounts of an anti-ErbB2 antibody and a cardioprotectant to the patient.

BACKGROUND OF THE INVENTION

Proto-oncogenes that encode growth factors and growth factor receptors have been identified to play important roles in the pathogenesis of various human malignancies, including breast cancer. It has been found that the human ErbB2 gene (erbB2, also known as her2, or c-erbB-2), which encodes a 185-kd transmembrane glycoprotein receptor ($p185^{HER2}$) related to the epidermal growth factor receptor (EGFR), is overexpressed in about 25% to 30% of human breast cancer (Slamon et al., *Science* 235:177-182 [1987]; Slamon et al., *Science* 244:707-712 [1989]).

Several lines of evidence support a direct role for ErbB2 in the pathogenesis and clinical aggressiveness of ErbB2-overexpressing tumors. The introduction of ErbB2 into non-neoplastic cells has been shown to cause their malignant transformation (Hudziak et al., *Proc. Natl. Acad. Sci. USA* 84:7159-7163 [1987]; DiFiore et al., *Science* 237:178-182 [1987]). Transgenic mice that express HER2 were found to develop mammary tumors (Guy et al., *Proc. Natl. Acad. Sci. USA* 89:10578-10582 [1992]).

Antibodies directed against human erbB2 protein products and proteins encoded by the rat equivalent of the erbB2 gene (neu) have been described. Drebin et al., *Cell* 41:695-706 (1985) refer to an IgG2a monoclonal antibody which is directed against the rat neu gene product. This antibody called 7.16.4 causes down-modulation of cell surface p185 expression on B104-1-1 cells (NIH-3T3 cells transfected with the neu proto-oncogene) and inhibits colony formation of these cells. In Drebin et al. *PNAS* (*USA*) 83:9129-9133 (1986), the 7.16.4 antibody was shown to inhibit the tumorigenic growth of neu-transformed NIH-3T3 cells as well as rat neuroblastoma cells (from which the neu oncogene was initially isolated) implanted into nude mice. Drebin et al. in *Oncogene* 2:387-394 (1988) discuss the production of a panel of antibodies against the rat neu gene product. All of the antibodies were found to exert a cytostatic effect on the growth of neu-transformed cells suspended in soft agar. Antibodies of the IgM, IgG2a and IgG2b isotypes were able to mediate significant in vitro lysis of neu-transformed cells in the presence of complement, whereas none of the antibodies were able to mediate high levels of antibody-dependent cellular cytotoxicity (ADCC) of the neu-transformed cells. Drebin et al. *Oncogene* 2:273-277 (1988) report that mixtures of antibodies reactive with two distinct regions on the p185 molecule result in synergistic anti-tumor effects on neu-transformed NIH-3T3 cells implanted into nude mice. Biological effects of anti-neu antibodies are reviewed in Myers et al., *Meth. Enzym.* 198:277-290 (1991). See also WO94/22478 published Oct. 13, 1994.

Hudziak et al., *Mol. Cell. Biol.* 9(3):1165-1172 (1989) describe the generation of a panel of anti-ErbB2 antibodies which were characterized using the human breast tumor cell line SKBR3. Relative cell proliferation of the SKBR3 cells following exposure to the antibodies was determined by crystal violet staining of the monolayers after 72 hours. Using this assay, maximum inhibition was obtained with the antibody called 4D5 which inhibited cellular proliferation by 56%. Other antibodies in the panel, including 7C2 and 7F3, reduced cellular proliferation to a lesser extent in this assay. Hudziak et al. conclude that the effect of the 4D5 antibody on SKBR3 cells was cytostatic rather than cytotoxic, since SKBR3 cells resumed growth at a nearly normal rate following removal of the antibody from the medium. The antibody 4D5 was further found to sensitize $p185^{erbB2}$-overexpressing breast tumor cell lines to the cytotoxic effects of TNF-α. See also WO89/06692 published Jul. 27, 1989. The anti-ErbB2 antibodies discussed in Hudziak et al. are further characterized in Fendly et al. *Cancer Research* 50:1550-1558 (1990); Kotts et al. In Vitro 26(3):59A (1990); Sarup et al. *Growth Regulation* 1:72-82 (1991); Shepard et al. *J. Clin. Immunol.* 11(3):117-127 (1991); Kumar et al. *Mol. Cell. Biol.* 11(2):979-986 (1991); Lewis et al. *Cancer Immunol. Immunother.* 37:255-263 (1993); Pietras et al. *Oncogene* 9:1829-1838 (1994); Vitetta et al. *Cancer Research* 54:5301-5309 (1994); Sliwkowski et al. *J. Biol. Chem.* 269(20):14661-14665 (1994); Scott et al. *J. Biol. Chem.* 266:14300-5 (1991); and D'souza et al. *Proc. Natl. Acad. Sci.* 91:7202-7206 (1994).

Tagliabue et al. *Int. J. Cancer* 47:933-937 (1991) describe two antibodies which were selected for their reactivity on the lung adenocarcinoma cell line (Calu-3) which overexpresses ErbB2. One of the antibodies, called MGR3, was found to internalize, induce phosphorylation of ErbB2, and inhibit tumor cell growth in vitro.

McKenzie et al. *Oncogene* 4:543-548 (1989) generated a panel of anti-ErbB2 antibodies with varying epitope specificities, including the antibody designated TA1. This TA1 antibody was found to induce accelerated endocytosis of ErbB2 (see Maier et al. *Cancer Res.* 51:5361-5369 [1991]). Bacus et al. *Molecular Carcinogenesis* 3:350-362 (1990) reported that the TA1 antibody induced maturation of the breast cancer cell lines AU-565 (which overexpresses the erbB2 gene) and MCF-7 (which does not). Inhibition of growth and acquisition of a mature phenotype in these cells was found to be associated with reduced levels of ErbB2 receptor at the cell surface and transient increased levels in the cytoplasm.

Stancovski et al. *PNAS* (*USA*) 88:8691-8695 (1991) generated a panel of anti-ErbB2 antibodies, injected them i.p. into nude mice and evaluated their effect on tumor growth of